

# Biocidal Surfaces May be an Important Adjunct in the Arsenal of Measures to Fight Healthcare-Acquired Infections

**Gadi Borkow\***

Cupron Scientific, Herzeliya, Israel

**\*Corresponding author:** Gadi Borkow, Cupron Scientific, Herzeliya, Israel, E-mail: gadi@cupron.com

## Editorial

Volume 2 Issue 2

**Received Date:** August 10, 2018


**Published Date:** August 14, 2018

## Editorial

Healthcare-acquired infections (HAIs) pose a critical threat to patients and have become a significant medical concern, especially as many microorganisms have developed high resistance to the existent antibiotics armory. Even in countries and settings where extensive infection control measures are routinely implemented, the risk of contracting HAIs by all types of microorganisms is very high. It is assessed that millions of people worldwide acquire a HAI each year. These infections contribute significantly to morbidity, mortality and hospitalization costs. For example, in the USA it is estimated that HAIs cause or contribute to ~100,000 deaths and add annually ~\$10 billion in additional healthcare expenses [1-3]. The risk of contracting a HAI in a clinical setting is extremely high, especially in developing countries. For example, the HAI rate reported for intensive care units in Morocco was 50%, in China 26.8%, in Turkey 29.1%, in Argentina 27%, in Mexico 24.4%, in India - 33.3%, in Jordan between 14.2 to 73.5%, and in Brazil between 8.3% to 60.8% [4-13].

In order to reduce HAI rates, aggressive measures have been implemented over the years by many medical entities. These include monitoring and isolation of infected patients, improved disinfection and hygiene regimens, increased number of nurses and infection control personnel, increased healthcare staff education, use of disposable equipment, and use of aggressive antibiotic control programs. These measures have resulted in significantly lower HAI rates; however, no elimination of HAI has been achieved even in settings where these infection control measures were rigorously implemented (Figure 1), and in many medical centers the HAI rates are still unacceptably high. It is clear that the current modalities to eliminate HAIs are not sufficient and

additional ways to fight HAI need to be taken into consideration. What are we missing and what else can we do?



**Figure 1:** Although rigorous infection control measures are being implemented it clear that the current modalities eliminate HAIs are not sufficient.

There is increasing evidence that contaminated non-intrusive soft and hard surfaces located in the clinical surroundings serve as a reservoir for nosocomial pathogens, which may then be transmitted to patients through contact and aerosolized particles (reviewed in [14]). The transmission may be direct or indirect. For example, patients may contaminate themselves directly by touching contaminated surfaces. Hospital personnel, even when using protective equipment such as gloves, may contaminate themselves by touching contaminated solid surfaces or textiles and then indirectly transfer the microorganisms to the patients. There is a clear correlation between the environmental bioburden present in a clinical setting and the risk of patients acquiring an infection [15-21]. Most common nosocomial

pathogens may persist on surfaces for months and can be a continuous source of microbial transmission [22,23].

Improved hygiene and better environmental decontamination contribute to control of epidemic and endemic transmission of many microorganisms, including antibiotic resistant and/or spore-forming bacteria, such as vancomycin-resistant enterococci, methicillin-resistant *Staphylococcus aureus*, and *Clostridium difficile* [24,25]. However, cleaning and disinfection do not always eradicate pathogens from surfaces [18,20]. Furthermore, the hospital environment is complex and often difficult to clean and the use of a cleaning agent that is not effective against the target organism can spread pathogens to other surfaces [14]. Only some of all hospital surfaces are routinely cleaned and enhanced surface cleaning practices in healthcare settings are difficult to sustain. Other approaches to eliminate environmental contamination, such as the use of ultraviolet irradiation and chemical fumigation, have significant limitations, for instance the need to remove the room occupants, and present safety risks [25].

A relatively new approach to overcoming the limitations of cleaning and disinfecting, and serving as an adjunct measure in the fight against environmental contamination and HAIs, is the use of self-disinfecting biocidal innate surfaces, especially those surrounding the patients [26-28]. The use in medical environments of surfaces containing copper or copper oxide has been found to reduce bioburden and the transmission of nosocomial pathogens [29-33]. For example, use of copper alloys surfaces in different clinical settings resulted in the reduction of bioburden from 58% to 98% as compared to matched controls. Furthermore, in a multi-center trial in which patients were randomly placed in rooms with or without copper alloy surfaces, there was a 58% reduction ( $P=0.013$ ) in the rate of HAI [30]. Methicillin-resistant *Staphylococcus aureus* (MRSA)/Vancomycin-resistant enterococci (VRE) patient colonization was found to be significantly lower by 42% in the copper arm of the study than in the control arm ( $P=0.02$ ). More recently, the use of linens, such as sheets, patient robes, patient pyjamas, and towels, containing copper oxide particles, has been found to reduce HAIs in several different clinical settings [34-37].

The use of self-disinfecting surfaces does not require special maintenance or personnel training, change in the routine work of the ward personnel and in the many measures already being taken to fight HAIs. Thus it may be an important adjunct to the arsenal of measures to fight Healthcare-Acquired Infections.

## References

1. Klevens RM, Edwards JR, Richards CL, Horan TC, Gaynes RP, et al. (2007) Estimating health care-associated infections and deaths in U.S. hospitals, 2002. *Public Health Rep* 122(2): 160-166.
2. Reed D, Kemmerly SA (2009) Infection control and prevention: a review of hospital-acquired infections and the economic implications. *Ochsner J* 9(1): 27-31.
3. Zimlichman E, Henderson D, Tamir O, Franz C, Song P, et al. (2013) Health care associated infections: a meta-analysis of costs and financial impact on the US health care system. *JAMA Intern Med* 173(22): 2039-2046.
4. Jroundi I, Khoudri I, Azzouzi A, Zeggwagh AA, Benbrahim NF, et al. (2007) Prevalence of hospital-acquired infection in a Moroccan university hospital. *Am J Infect Control* 35(6): 412-416.
5. Ding JG, Sun QF, Li KC, Zheng MH, Miao XH, et al. (2009) Retrospective analysis of nosocomial infections in the intensive care unit of a tertiary hospital in China during 2003 and 2007. *BMC Infect Dis* 9: 115-120.
6. Inan D, Saba R, Yalcin AN, Yilmaz M, Ongut G, et al. (2006) Device-associated nosocomial infection rates in Turkish medical-surgical intensive care units. *Infect Control Hosp Epidemiol* 27(4): 343-348.
7. Rosenthal VD, Guzman S, Orellano PW (2003) Nosocomial infections in medical-surgical intensive care units in Argentina: attributable mortality and length of stay. *Am J Infect Control* 31(5): 291-295.
8. Ramirez Barba EJ, Rosenthal VD, Higuera F, Oropeza MS, Hernandez HT, et al. (2006) Device-associated nosocomial infection rates in intensive care units in four Mexican public hospitals. *Am J Infect Control* 34(4): 244-247.
9. Agarwal R, Gupta D, Ray P, Aggarwal AN, Jindal SK (2006) Epidemiology, risk factors and outcome of nosocomial infections in a Respiratory Intensive Care Unit in North India. *J Infect* 53(2): 98-105.
10. Khuri-Bulos NA, Shennak M, Agabi S, Saleh S, Al Rawashdeh S, et al. (1999) Nosocomial infections in the intensive care units at a university hospital in a developing country: comparison with National

- Nosocomial Infections Surveillance intensive care unit rates. *Am J Infect Control* 27(6): 547-552.
11. Abramczyk ML, Carvalho WB, Carvalho ES, Medeiros EA (2003) Nosocomial infection in a pediatric intensive care unit in a developing country. *Braz J Infect Dis* 7(6): 375-380.
  12. Moura ME, Campelo SM, de Brito FC, Batista OM, de Araujo TM, et al. (2007) Nosocomial infection: study of prevalence at a public teaching hospital. *Rev Bras Enferm* 60(4): 416-421.
  13. Salomao R, Rosenthal VD, Grimberg G, Nouer S, Blecher S, et al. (2008) Device-associated infection rates in intensive care units of Brazilian hospitals: findings of the International Nosocomial Infection Control Consortium. *Rev Panam Salud Publica* 24(3): 195-202.
  14. Otter JA, Yezli S, French GL (2014) The role of contaminated surfaces in the transmission of nosocomial pathogens. In: *Use of biocidal surfaces for reduction of healthcare acquired infections*. (Borkow G. Ed.) Chapter 3: 27-58.
  15. Hota B (2004) Contamination, disinfection, and cross-colonization: are hospital surfaces reservoirs for nosocomial infection? *Clin Infect Dis* 39(8): 1182-1189.
  16. Bhalla A, Pultz NJ, Gries DM, Ray AJ, Eckstein EC, et al. (2004) Acquisition of nosocomial pathogens on hands after contact with environmental surfaces near hospitalized patients. *Infect Control Hosp Epidemiol* 25(2): 164-167.
  17. Boyce JM, Potter-Bynoe G, Chenevert C, King T (1997) Environmental contamination due to methicillin-resistant *Staphylococcus aureus*: possible infection control implications. *Infect Control Hosp Epidemiol* 18(9): 622-627.
  18. Eckstein BC, Adams DA, Eckstein EC, Rao A, Sethi AK, et al. (2007) Reduction of *Clostridium Difficile* and vancomycin-resistant *Enterococcus* contamination of environmental surfaces after an intervention to improve cleaning methods. *BMC Infect Dis* 7: 61.
  19. Goodman ER, Platt R, Bass R, Onderdonk AB, Yokoe DS, et al. (2008) Impact of an environmental cleaning intervention on the presence of methicillin-resistant *Staphylococcus aureus* and vancomycin-resistant enterococci on surfaces in intensive care unit rooms. *Infect Control Hosp Epidemiol* 29(7): 593-599.
  20. Manian FA, Griesenauer S, Senkel D, Setzer JM, Doll SA, et al. (2011) Isolation of *Acinetobacter baumannii* complex and methicillin-resistant *Staphylococcus aureus* from hospital rooms following terminal cleaning and disinfection: can we do better? *Infect Control Hosp Epidemiol* 32(7): 667-672.
  21. Sabino R, Sampaio P, Carneiro C, Rosado L, Pais C (2011) Isolates from hospital environments are the most virulent of the *Candida parapsilosis* complex. *BMC Microbiol* 11(1): 180.
  22. Kramer A, Schwebke I, Kampf G (2006) How long do nosocomial pathogens persist on inanimate surfaces? A systematic review. *BMC Infect Dis* 6: 130-138.
  23. Kramer A, Assadian O (2014) Survival of microorganisms on inanimate surfaces. In: *Use of biocidal surfaces for reduction of healthcare acquired infections*. Borkow G (Edr.), 7-26.
  24. Hayden MK, Bonten MJ, Blom DW, Lyle EA, van de Vijver DA, et al. (2006) Reduction in acquisition of vancomycin-resistant enterococcus after enforcement of routine environmental cleaning measures. *Clin Infect Dis* 42(11): 1552-1560.
  25. Byrns G (2014) Alternative room disinfection modalities - pros and cons. In: *Use of biocidal surfaces for reduction of healthcare acquired infections* 9: 187-207.
  26. Noyce JO, Michels H, Keevil CW (2006) Potential use of copper surfaces to reduce survival of epidemic methicillin-resistant *Staphylococcus aureus* in the healthcare environment. *J Hosp Infect* 63(3): 289-297.
  27. Borkow G, Gabbay J (2008) Biocidal textiles can help fight nosocomial infections. *Med Hypotheses* 70(5): 990-994.
  28. Borkow G, Monk AB (2012) Fighting nosocomial infections with biocidal nonintrusive hard and soft surfaces. *World J Clin Infect Dis* 12(4): 77-90.
  29. Mikolay A, Huggett S, Tikana L, Grass G, Braun J, et al. (2010) Survival of bacteria on metallic copper surfaces in a hospital trial. *Appl Microbiol Biotechnol* 87(5): 1875-1879.

30. Salgado CD, Sepkowitz KA, John JF, Cantey JR, Attaway HH, et al. (2013) Copper surfaces reduce the rate of healthcare-acquired infections in the intensive care unit. *Infect Control Hosp Epidemiol* 34(5): 479-486.
31. Schmidt MG, Attaway HH, Sharpe PA, John J, Sepkowitz KA, et al. (2012) Sustained reduction of microbial burden on common hospital surfaces through introduction of copper. *J Clin Microbiol* 50(7): 2217-2223.
32. Monk AB, Kanmukhla V, Trinder K, Borkow G (2014) Potent bactericidal efficacy of copper oxide impregnated non-porous solid surfaces. *BMC Microbiol* 14(1): 57.
33. Sifri CD, Burke GH, Enfield KB (2016) Reduced health care-associated infections in an acute care community hospital using a combination of self-disinfecting copper-impregnated composite hard surfaces and linens. *Am J Infect Control* 44(12): 1565-1571.
34. Lazary A, Weinberg I, Vatine JJ, Jefidoff A, Bardenstein R, et al. (2014) Reduction of healthcare-associated infections in a long-term care brain injury ward by replacing regular linens with biocidal copper oxide impregnated linens. *Int J Infect Dis* 24: 23-29.
35. Marcus EL, Yosef H, Borkow G, Caine Y, Sasson A, et al. (2017) Reduction of health care-associated infection indicators by copper oxide-impregnated textiles: Crossover, double-blind controlled study in chronic ventilator-dependent patients. *Am J Infect Control* 45(4): 401-403.
36. Burke GH, Butler Jp (2018) Analysis of the role of copper impregnated composite hard surfaces, bed linens and patient gowns in reducing *Clostridium difficile* and multi drug resistant organism healthcare associated infection rates using a multi facility analysis within a healthcare system. *International Journal of Infection Control* 14(1): 1-8.
37. Butler JP (2018) Effect of copper-impregnated composite bed linens and patient gowns on healthcare-associated infection rates in six hospitals. *J Hosp Infect.* Epub ahead of print.

